TEST NAME/ PSYCHE CODE	COLLECTION GUIDELINES	REFERENCE VALUES	ROOM TEMP.	REFRIGERATED	FROZEN	Clinical Utility
URIC ACID, SERUM	1 ml SERUM. FASTING PREFERRED.	2.2-7.0 mg/dL		48 hours		The uric acid blood test is used to detect high levels of this compound in the blood in order to help diagnose
CODE: URIC						gout.
CPT: 84550	SCHEDULE: DAILY					
URINALYSIS (MICROSCOPIC PREFORMED IF INDICATED)	MIDSTREAM SPECIMEN PREFERRED IN CLEAN CONTAINER. SUBMIT AT LEAST 10 ml URINE. REFRIGERATE	INTERPRETIVE REPORT		48 hours		The urinalysis is used as a screening and/or diagnostic tool because it can help detect substances or cellular material in the urine associated with different metabolic and kidney disorders.
CODE: UA						
CPT: 81001	SCHEDULE: DAILY					
URINALYSIS WITH MICROSCOPIC	MIDSTREAM SPECIMEN PREFERRED IN CLEAN CONTAINER. SUBMIT AT LEAST 10 ml URINE. REFRIGERATE.	INTERPRETIVE REPORT		48 hours		The urinalysis is used as a screening and/or diagnostic tool because it can help detect substances or cellular material in the urine associated with different metabolic and kidney disorders.
CODE: UAWM						
CPT: 81003	SCHEDULE: DAILY					
				48 hours		The urine culture test is used to diagnose a urinary tract infection (UTI).
URINE CULTURE IF INDICATED CODE: CULTURE IF INDICATED *NOTE ON REQUISITION	MUST BE WRITTEN ON REQUISITION. CULTURE PERFORMED IF UA SHOWS ONE OF MORE OF THE FOLLOWING: 1. LEUKOCYTE ESTERASE OF 1+ OR GREATER 2. POSTIVE NITRATES 3. 6-10 WBC/HPF OR GREATER 4. 1+ OR GREATER BACTERIA/YEAST	INTERPRETIVE REPORT				